

SCIENTIFIC NOTE

Notes on the status of the Eurasian moths *Noctua pronuba* and *Noctua comes* (Lepidoptera: Noctuidae) on Vancouver Island, British ColumbiaCLAUDIA R. COPLEY^{1,2} and ROBERT A. CANNINGS²

Two Eurasian cutworm moths (Lepidoptera: Noctuidae), *Noctua pronuba* (Linnaeus) and *Noctua comes* (Hübner), both accidentally introduced to North America, are now sympatric in southwestern British Columbia. The former was first recorded in Nova Scotia, the latter in British Columbia. This paper reports the occurrence of both species for the first time on Vancouver Island. They are the only species of the genus *Noctua* known in North America (Lafontaine 1998).

Noctua pronuba (the large yellow underwing) was first reported in North America in Nova Scotia in 1979 (Neil 1981). It is now known from every Canadian province and Nunavut (Troubridge and Lafontaine 2005) and, in the USA, from Maine (Wright 1987) to Louisiana (Brou 1997) and California (Powell 2002). It was first recorded in BC in 2002 (CNC [Canadian National Collection of Insects and Arachnids, Ottawa] data) and is now abundant on eastern Vancouver Island as far north as Sayward (RBCM [Royal British Columbia Museum, Victoria] data) and areas of the lower Fraser River Valley (K. Needham, pers. comm.). We have yet to hear of any records from the BC Interior.

Noctua comes (the lesser yellow underwing) was first recorded in Canada in Burnaby, BC in August 1982 (Neil 1984) although a specimen in the Spencer Entomological Museum, UBC (University of British Columbia, Vancouver) was collected in Vancouver in July 1982. This species was first recorded on Vancouver Island in Victoria in 1990 (PFC [Pacific Forestry Centre, Victoria] data) and is now abundant in suburban habitats there. Elsewhere in BC *N.*

comes has been found in the Okanagan Valley and Lillooet and south to Oregon (Lafontaine 1998, J. Troubridge, pers. comm.).

Noctua pronuba has a wingspan of 50–60 mm and a diagnostic orange-yellow hindwing with a broad black border. Images of adults and larvae are in Wright (1987), Lafontaine (1998), and Neil and Specht (1987). *Noctua comes* is similar but normally has a black mark near the centre of the orange of the hindwing; genitalia differences distinguish the two species unequivocally (Lafontaine 1998).

Although *N. pronuba* is known to be migratory and a very strong flier (Passoa and Hollingsworth 1996), its spread may have been facilitated by human activity. It has a wide range of host plants, many of which are part of the horticultural trade, food-crop industry, or are widespread weeds. Host plant genera include: *Holcus* (J. Tatum, pers. comm.), *Poa* and other grasses (Wright and Neil, 1983), *Atriplex*, *Chrysanthemum*, *Dianthus*, *Fragaria*, *Free-sia*, *Gladiolus*, *Myosotis*, *Polygonum*, *Primula*, *Ribes*, *Stellaria*, *Taraxacum* and *Viola*. Larvae also eat various common food crops (Passoa and Hollingsworth 1996, B. Duncan, pers. comm.). *Noctua comes* has been recorded on *Conium*, *Cornus*, *Potentilla*, *Calendula*, *Cardamine*, *Cirsium*, *Digitalis*, *Fragaria*, *Myosotis*, *Plantago*, *Primula* but, most often, on *Rumex crispus* (J. Tatum, pers. comm.) as well as tobacco and grapes (Sannino and Espinosa 1999) and *Crataegus* (Ward 2003).

Life-history details of *N. pronuba* are reported in Singh and Kevan (1965),

¹ Corresponding author, email: ccopley@royalbcmuseum.bc.ca

² Royal British Columbia Museum, 675 Belleville Street, Victoria, BC, V8W 9W2

Wright and Neil (1983), Morris (1987), and Passoa and Hollingsworth (1996). In British Columbia, *N. pronuba* will likely exhibit the univoltine life history typical of European and eastern North American populations.

Each female lays up to 2000 eggs on leaf undersides (Morris 1987) or non-host substrates (B. Duncan, pers. comm.). Larvae feed on foliage, crowns and roots of hosts; immature larvae usually overwinter (Morris 1987), but in coastal BC mature larvae also do so (B. Duncan, pers. comm.). Mature larvae pupate in the soil in the spring (Passoa and Hollingsworth 1996). Tachinid flies parasitize *N. pronuba* in eastern North America (J. Troubridge, pers. comm.), but have not been recorded in BC on *N. comes* or *N. pronuba* (J. Tatum, pers. comm.). However, *Trichogramma* wasps parasitize egg masses in the province (B. Duncan, pers. comm.).

As it is a strong, migratory flier (Passoa and Hollingsworth 1996), can endure very cold winters (Wright and Neil 1983), and feeds on a wide array of plants associated

with humans, *N. pronuba* will probably colonize all of BC. Adults may lay eggs on non-plant substrates (B. Duncan, pers. comm) or hide during the day in objects around human habitation (although they fly readily when disturbed), making them good candidates for transport by vehicles.

Because of its growing abundance in coastal BC, we believe that *N. pronuba* may become an economic pest, although in the long-term, populations will likely be moderated by increasing parasitism. *Noctua comes* will probably have a similar future.

We thank Jane Seed, Karen Needham and Chris Borkent for data from PFC, UBC and CNC collections, respectively and the following colleagues for their permission to use their personal communications. Jim Troubridge (Agriculture and Agri-Food Canada, Ottawa, ON) and Karen Needham (UBC, Vancouver, BC) provided information on distribution. Jeremy Tatum (University of Victoria, Victoria, BC) and Bob Duncan (PFC, Victoria, BC) commented on food plants and life history.

REFERENCES

- Brou, V. 1997. A Gulf Coast record of the European cutworm, *Noctua pronuba* (L.). *News of the Southern Lepidopterists' Society* 19: 3.
- Lafontaine, J.D. 1998. Noctuoidea, Noctuidae (part). In Dominick, R.B. *et al.* The Moths of America North of Mexico, Fascicle 27.3. The Wedge Entomological Research Foundation, Washington, DC.
- Morris, R.F. 1987. Note on occurrence of the large yellow underwing moth, *Noctua pronuba* (Linnaeus) (Lepidoptera: Noctuidae), in Newfoundland. *The Canadian Entomologist* 119: 403-404.
- Neil, K.A. 1981. The occurrence of *Noctua pronuba*, (Noctuidae) in Nova Scotia: a new North America record. *Journal of the Lepidopterist's Society* 35: 248.
- Neil, K.A. 1984. *Noctua comes*, a noctuid new to North America (Lepidoptera: Noctuidae: Noctuinae). *The Canadian Entomologist* 116: 479-480.
- Neil, K.A. and H.B. Specht. 1987. Sixth-instar larvae of *Noctua pronuba* (L.) (Lepidoptera: Noctuidae). *The Canadian Entomologist* 119: 209-214.
- Passoa, S. and C.S. Hollingsworth. 1996. Distribution, identification and rate of spread of *Noctua pronuba* (Lepidoptera: Noctuidae) in the northeastern United States. *Entomological News* 107: 151-160.
- Powell J.A. 2002. *Noctua pronuba* reaches the Pacific Coast. *News of the Lepidopterists' Society* 44: 120.
- Sannino, L. and B. Espinosa. 1999. On the morphology of *Noctua comes* (Lepidoptera: Noctuidae). *Il Tabacco* 7: 35-43.
- Singh, M.P. and D.K. Kevan. 1965. Notes on three common British species of agrotid moth. *Entomologists Record and Journal of Variation* 68: 233-235.
- Troubridge, J.T. and J.D. Lafontaine. 2005. The Moths of Canada. Part 1, Noctuoidea of Western Canada and Part 2, Noctuoidea of Eastern Canada. Canadian Biodiversity Information Facility, Government of Canada, Ottawa, ON. http://www.cbif.gc.ca/spp_pages/misc_moths/phps/mothindex_e.php.
- Ward, J. 2003. Moths of Northamptonshire. Northamptonshire Moth Group. <http://www.northamptonshirewildlife.co.uk/nmoths/nmoths.htm>
- Wright, B. 1987. The European yellow underwing, *Noctua pronuba* (L.) (Lepidoptera: Noctuidae) in the Atlantic provinces (Canada) and the State of Maine (USA). *The Canadian Entomologist* 119: 993-997.
- Wright, B. and K.A. Neil. 1983. *Noctua pronuba*, a European cutworm, established in Nova Scotia (Lepidoptera: Noctuidae). *The Canadian Entomologist* 115: 1047-1048.